

Notes on the minimal longitudinal dc conductivity of perfect bilayer graphene

József Cserti¹¹*Department of Physics of Complex Systems, Eötvös University
H-1117 Budapest, Pázmány Péter sétány 1/A, Hungary*

We calculated the minimal longitudinal conductivity in perfect single and bilayer graphene by extending the two methods developed for Dirac fermion gas by A. W. W. Ludwig et al. in Phys. Rev. B **50**, 7526 (1994). Using the Kubo formula which was originally applied for spintronic systems we obtain $\sigma_{xx}^{min} = (J\pi/2) e^2/h$ while from the other formula used in the above mentioned work we find $\bar{\sigma}_{xx}^{min} = (4J/\pi) e^2/h$, where $J = 1$ for single layer and $J = 2$ for bilayer graphene. The two universal values are different although they are numerically close to each other. Our two results are in the same order of magnitude as that of experiments and for single layer case one of our result agrees many earlier theoretical predictions. However, for bilayer graphene only two studies are known with predictions for the minimal conductivity different from our calculated values. Similarly to the single layer case, the physical origin of the minimal conductivity in bilayer graphene is also rooted back to the intrinsic disorder induced by the Zitterbewegung which is related to the trembling motion of the electron.

PACS numbers: 81.05.Uw, 73.23.Ad, 72.10.Bg, 73.43.Cd

Unusual and remarkable transport properties of graphene (single or stacks of atomic layer of graphite) have been proved by recent experiments^{1,2}. Besides the unconventional quantum Hall effect, many new phenomena have been predicted and studied in graphene such as the Klein paradox³, the specular Andreev reflection^{4,5}, the Josephson effect⁶, a new electric field effect⁷, the photon-assisted electron transport⁸, composite Dirac fermions⁹, quantum dots¹⁰, the n-p junction¹¹, the fractional quantum Hall effect¹², the spin-orbit gap¹³.

In this work we study the longitudinal conductivity of graphenes which, according to experiments, takes the minimum values of orders of e^2/h . This is an intrinsic property in the sense that it persists even in perfect (impurity free) carbon honeycomb lattice. Such a peculiar behavior can be related to the excitation spectrum of single layer graphene described well by a Dirac like dispersion relation (Dirac cones) of two-dimensional massless Dirac fermions¹⁴.

Much theoretical efforts have been devoted to explain quantitatively the observed minimal longitudinal conductivity. However, at the theoretical side different predictions exist for the value of the minimal conductivity σ_{xx}^{min} even in perfect single layer graphene. Actually, long before the experimental evidence of the minimal conductivity, it has already been considered theoretically^{15,16,17} and Ludwig et al. found different values using two different approaches¹⁸. Similarly, in many recent works^{19,20,21,22} it has been found that $\sigma_{xx}^{min} = (4/\pi) e^2/h$, while Ziegler predicted²³ $\sigma_{xx}^{min} = \pi e^2/h$, Falkovsky and Varlamov²⁴ obtained $\sigma_{xx}^{min} = (\pi/2) e^2/h$, and Nomura and MacDonald obtain from numerical calculations of the Kubo formula $\sigma_{xx}^{min} = (4/\pi) e^2/h$ for short range scattering case and $\sigma_{xx}^{min} = 4 e^2/h$ for Coulomb scattering case²⁵. Recently, Gusynin and Sharapov have derived analytical results for the ac and dc conductivity of Dirac fermions in graphene²⁶ which could be used for studying the minimal conductivity.

For bilayer graphene first studied experimentally²⁷ by Novoselov et al. and theoretically²⁸ by McCann and Fal'ko, the situation is not better regarding the minimal conductivity. Much less theoretical works considered the minimal conductivity in bilayer graphene. Recently, Koshino and Ando have investigated the transport in bilayer graphene in self-consistent Born approximation²⁹ and they found that in the strong-disorder regime $\sigma_{xx}^{min} = (8/\pi) e^2/h$, while in the weak-disorder regime $\bar{\sigma}_{xx}^{min} = (24/\pi) e^2/h$ which is six times larger than in single layer graphene. Similarly, Katsnelson has also calculated the minimal conductivity in bilayers using the Landauer approach^{29,30} and he obtained a different value $\sigma_{xx}^{min} = 2 e^2/h$.

For massless Dirac fermion (corresponding to the single layer graphene) Ludwig et al. calculated the conductivity¹⁸ using two definitions for that, the one is the Kubo formula (σ_{xx}^{min}) and the other ($\bar{\sigma}_{xx}^{min}$) is a definition often used in the sigma model literature³¹. The two definitions yield two different results (although numerically they are close to each other) for the longitudinal conductivity of perfect single layer graphene, namely, $\sigma_{xx}^{min} = n_v n_s (\pi/8) e^2/h$ and $\bar{\sigma}_{xx}^{min} = n_v n_s (1/\pi) e^2/h$, respectively. Here the factor $n_v = 2$ and $n_s = 2$ correspond to the two valleys and the electron spin, respectively (for details see references before).

As can be seen from the above reviewed literature, several predictions of the exact value of the minimal conductivity even in perfect single and bilayer graphene exist although they are consistent at least in order of magnitude. In this work we extend Ludwig et al. approach to calculate the longitudinal conductivity for bilayer graphene. We find that, similarly to the case of single layer graphene, the two definitions of the conductivity used in Ref. 18 yield also different results for perfect bilayer graphene. More interestingly, we find that the conductivity obtained from both approaches is *doubled* compared to that for single layer graphene. Thus,

our results show that obtaining the exact value of the minimal conductivity is a rather subtle task and it needs further investigation.

Regarding the technical details of our calculations we note that when we used the Kubo formula, instead of following the approach of Ludwig et al. we applied an *equivalent* method developed by Bernevig³² for spintronic systems. In this method the conductivity is calculated in ‘bubble’ approximation using the one-particle non-interacting Green’s function for finite temperature. This approach results in the same conductivity for single layer graphene as that obtained from Ludwig et al. method and, as can be seen below, it is more convenient to extend to bilayer graphene.

In order to treat the single and bilayer graphene simultaneously we start from the Hamiltonian given in a unified form as^{1,28}

$$H_J = g \begin{pmatrix} 0 & (p_x - ip_y)^J \\ (p_x + ip_y)^J & 0 \end{pmatrix}, \quad (1)$$

where $J = 1$ for single and $J = 2$ for bilayer graphene, and g is a constant depending on J . In what follows, it is more convenient to use the following equivalent form of Eq. (1)

$$H_J = \Omega(\mathbf{p}) \boldsymbol{\sigma}, \quad (2)$$

where $\boldsymbol{\sigma} = (\sigma_x, \sigma_y)$ are the Pauli matrices representing the ‘pseudo spin’ acting on the spinor states with components corresponding to the wave function’s amplitudes at the two non-equivalent basis atoms in the unit cell of the honeycomb lattice, $\Omega(\mathbf{p}) = gp^J(\cos J\varphi, \sin J\varphi)$ and φ is the polar angle of \mathbf{p} , i.e., $\mathbf{p} = p(\cos \varphi, \sin \varphi)$ with $p = |\mathbf{p}|$. The eigenstates of the Hamiltonian H_J with plane wave solutions $e^{i\mathbf{k}\mathbf{r}}$ have eigenvalues

$$E_{\pm}(\mathbf{k}) = \pm\Omega(\hbar\mathbf{k}), \quad (3)$$

where $\Omega = \sqrt{\Omega^2(\hbar\mathbf{k})}$ is the magnitude of the vector $\Omega(\hbar\mathbf{k})$.

The Green’s function of a Hamiltonian H is defined by $\hat{G}(z) = (z - \hat{H})^{-1}$ and its position representation reads $G(\mathbf{r}, \mathbf{r}', z) = \langle \mathbf{r} | \hat{G}(z) | \mathbf{r}' \rangle$. Using the Hamiltonian (2) one finds

$$\begin{aligned} G(\mathbf{r}, \mathbf{r}', z) &= \langle \mathbf{r} | G(z) | \mathbf{r}' \rangle \\ &= \int \frac{d^2k}{(2\pi)^2} e^{i\mathbf{k}(\mathbf{r}-\mathbf{r}')} \frac{z + \Omega(\mathbf{k}) \boldsymbol{\sigma}}{z^2 - \Omega^2(\mathbf{k})}, \end{aligned} \quad (4)$$

where z is a complex number.

To calculate the conductivity from the Kubo formula we apply the result derived first by Bernevig³² for spin-

tronic systems using the Green’s function (4):

$$\sigma_{ij}(\omega) = -\frac{e^2}{\hbar} \lim_{\omega \rightarrow 0} \frac{\text{Im}\{\Pi_{ij}^{\text{ret}}(\omega)\}}{\omega}, \quad \text{where} \quad (5a)$$

$$\begin{aligned} \Pi_{ij}(\omega) &= \frac{2}{A} \sum_{\mathbf{k}} \frac{n_F(E_+) - n_F(E_-)}{\omega^2 - 4\Omega^2} \Omega^2 \\ &\times \left(i\omega \epsilon_{\alpha\beta\gamma} F_{i\alpha} F_{j\beta} \hat{\Omega}_{\gamma} - 2\Omega F_{i\alpha} F_{j\alpha} \right), \end{aligned} \quad (5b)$$

$$F_{i\alpha} = \frac{\partial \hat{\Omega}_{\alpha}}{\partial k_i}, \quad \hat{\Omega}_{\alpha} = \frac{\Omega_{\alpha}}{\Omega}, \quad (5c)$$

and $\Pi_{ij}^{\text{ret}}(\omega)$ is the retarded correlation function obtained by analytic continuation $\omega \rightarrow \omega + i\eta$ of function $\Pi_{ij}(\omega)$, $n_F(E) = 1/(e^{\beta(E-\mu)} + 1)$ is the Fermi function with Fermi energy μ , the indices take the values 1, 2, 3, and summation over repeated indices is assumed. Here A is the area of the two dimensional system, $\epsilon_{\alpha\beta\gamma}$ is the Levi-Civita symbol, and the two energy bands $E_{\pm}(\mathbf{k})$ are given by Eq. (3). This result is valid for ballistic systems (note that this result is slightly rewritten to make it more transparent and useful for further calculations).

We now evaluate the conductivity for undoped single and bilayer graphene at zero temperature using Eq. (5). The Fermi energy for undoped graphene is $\mu = 0$, i.e., the negative energy band is fully occupied and the positive one is empty at zero temperature. Thus, in Eq. (5b) $n_F(E_+) - n_F(E_-) = -1$ for all occupied states with Fermi wave number \mathbf{k} . For dc conductivity we need to find $\sigma(\omega)$ in the limit $\omega \rightarrow 0$. Therefore, the integration over \mathbf{k} can safely be extended to infinity since the contribution comes from the integrand close to $k \approx 0$ (see the denominator of $\Pi_{ij}(\omega)$). Using Eq. (5b) and $\sum_{\mathbf{k}} \rightarrow A \int d^2k/(2\pi)^2$ (no contribution comes from the term contains the Levi-Civita symbol $\epsilon_{\alpha\beta\gamma}$) we find

$$\Pi_{xx}^{\text{ret}}(\omega) = \lim_{\eta \rightarrow 0^+} \int_0^{\infty} \frac{dk}{2\pi} \frac{2g^3 J^2 k^{3J-1}}{(\omega + i\eta)^2 - 4g^2 k^{2J}}. \quad (6)$$

The imaginary part of the above integral can be evaluated analytically and it yields

$$\text{Im}\{\Pi_{xx}(\omega)\} = -\frac{J\omega}{16}. \quad (7)$$

Thus, from (5a) the longitudinal dc conductivity (per valley per spin) for single ($J = 1$) and bilayer ($J = 2$) graphene becomes:

$$\sigma_{xx} = \frac{J\pi}{8} \frac{e^2}{\hbar}. \quad (8)$$

Note that the conductivity obtained from the Kubo formula is universal depending only on the type of graphene but not the constant g .

We now apply the other definition used by Ludwig et al. for calculating the conductivity (see Eq. (55) in

Ref. 18) which reads as

$$\bar{\sigma}_{xx} = \frac{e^2}{h} \lim_{\eta \rightarrow 0^+} \eta^2 \int d^2r r^2 \text{Tr} [G(0, \mathbf{r}, i\eta) G(\mathbf{r}, 0, -i\eta)] \quad (9)$$

where the Green's function is given by Eq. (4). The conductivity $\bar{\sigma}_{xx}$ can be rewritten as^{18,31}

$$\bar{\sigma}_{xx} = -\frac{8e^2}{h} \lim_{\eta \rightarrow 0^+} \eta^2 \frac{\partial}{\partial q^2} \Big|_{q=0} K(\mathbf{q}, \eta), \quad (10a)$$

$$K(\mathbf{q}, \eta) = \frac{1}{2} \int d^2r e^{i\mathbf{q}\mathbf{r}} \text{Tr} [G(0, \mathbf{r}, i\eta) G(\mathbf{r}, 0, -i\eta)], \quad (10b)$$

and the trace is taken over the 'pseudo spin space'.

Using the Green's function given by Eq. (4) and performing the trace over the 'pseudo spin space' in (10b) we find

$$K(\mathbf{q}, \eta) = \int \frac{d^2k}{(2\pi)^2} \frac{\boldsymbol{\Omega}_+ \boldsymbol{\Omega}_- + \eta^2}{(\boldsymbol{\Omega}_+^2 + \eta^2)(\boldsymbol{\Omega}_-^2 + \eta^2)}, \quad (11a)$$

$$\boldsymbol{\Omega}_{\pm} = \boldsymbol{\Omega}(\mathbf{k} \pm \mathbf{q}/2). \quad (11b)$$

After a simple algebra the integrand I in Eq. (11a) can be calculated explicitly:

$$I = \frac{g^2 \left[(k^2 + q^2/4)^2 - (\mathbf{k}\mathbf{q})^2 \right]^{J/2} \cos(J\Phi) + \eta^2}{\Sigma_+ \Sigma_-} \quad (12a)$$

$$\Sigma_{\pm} = g^2 \left(k^2 + \frac{q^2}{4} \pm \mathbf{k}\mathbf{q} \right)^J + \eta^2, \quad (12b)$$

$$\cos \Phi = \frac{k^2 - q^2/4}{\sqrt{(k^2 + q^2/4)^2 - (\mathbf{k}\mathbf{q})^2}}. \quad (12c)$$

Here Φ is the angle between vector $\mathbf{k} + \mathbf{q}/2$ and $\mathbf{k} - \mathbf{q}/2$. Expanding I for small q we obtain

$$I \approx \frac{1}{g^2 k^{2J} + \eta^2} - q^2 g^2 J k^{2J-2} \frac{J (g^2 k^{2J} + 3\eta^2) - [g^2 (J+1) k^{2J} - (J-1) \eta^2] \cos(2\varphi - 2\gamma)}{4(g^2 k^{2J} + \eta^2)^3}, \quad (13)$$

where φ and γ are the polar angle of \mathbf{k} and \mathbf{q} , i.e., $\mathbf{k} = k(\cos \varphi, \sin \varphi)$ and $\mathbf{q} = q(\cos \gamma, \sin \gamma)$. As can be seen below γ will be dropped out in the final result.

Integrating the integrand I in (13) over \mathbf{k} and taking the derivation with respect to q^2 we have the following simple result:

$$\frac{\partial}{\partial q^2} \Big|_{q=0} K(\mathbf{q}, \eta) = -\frac{J}{8\pi\eta^2}, \quad (14)$$

where γ has been dropped out after the integration over φ . Similarly, the first term disappears in I since it is independent of q . Now using Eq. (10a) the conductivity (per valley per spin) becomes

$$\bar{\sigma}_{xx} = \frac{J}{\pi} \frac{e^2}{h}. \quad (15)$$

Discussion: We calculated the minimal conductivity for perfect single ($J = 1$) and bilayer ($J = 2$) graphene using the Kubo formula and the definition (9). Taking into account the two valleys ($n_v = 2$) and the electron spin ($n_s = 2$) we find from Eqs. (8) and (15) that the minimal conductivities take the universal values:

$$\sigma_{xx}^{min} = \frac{J\pi}{2} \frac{e^2}{h}, \quad (16a)$$

$$\bar{\sigma}_{xx}^{min} = \frac{4J}{\pi} \frac{e^2}{h}. \quad (16b)$$

Note that these results are independent of the constant g appeared in the Hamiltonian (1). Just like in the case of single layer graphene we also find different values for the conductivity when the two methods in Ref. 18 is extended to bilayer graphene. Moreover, the minimal conductivity in perfect bilayer graphene calculated from Eqs. (5) and (9) is doubled compared to that in single layer graphene. One can also see that for perfect single layer graphene our result (16b) agrees with that obtained in Refs. 19,20, 21,22.

As pointed out by Katsnelson^{21,30}, the reason for the different results (16a) and (16b) is related to the 'trembling' or oscillatory motion of the center of a free wave packet called *Zitterbewegung* which has recently been reconsidered in Ref. 33. The *Zitterbewegung* induces an intrinsic disorder in perfect graphenes which has indirectly been confirmed by Tworzydło et al.²² by calculating the Fano factor of the shot noise in single layer graphene, and they found that it has the same value (1/3) as in disordered metals.

In our calculation the energy dispersion of bilayer graphenes was assumed to be purely parabolic in \mathbf{k} (see Eq. (1)). Koshino and Ando studied the conductivity when the energy dispersion includes both \mathbf{k} -linear and \mathbf{k} -square terms²⁹ which results in a trigonally warped Fermi surface²⁸. To clarify the role of the \mathbf{k} -linear term it would

be interesting to extend our work along this line.

Similarly, extending the theoretical efforts to study multi-layer graphenes would be desirable in order to have better understanding of the unusual transport in graphenes. Recently, the electronic states in stacks of graphene layers have been studied³⁴. However, it is not clear whether in the case of trilayers the Hamiltonian can be described simple by Eq. (1) with $J = 3$.

The single and bilayer graphene are zero-gap semiconductors but this gap can open up due to impurities or the presence of interaction. Then, graphenes mapped to the massless Dirac fermion gas transform to random mass Dirac fermion systems. In this case, to calculate the con-

ductivity it is useful to start from the definition (9) as it was demonstrated by Ludwig et al.¹⁸. Recently, Gusynin et al. proposed theoretically that microwaves could be an experimental tool to understand the role of the impurities and the presence of interaction in graphene systems³⁵.

In summary, our work shows that the physical explanation of the existence of the minimal conductivity in graphene systems still remains a theoretical challenge in the future.

We gratefully acknowledge discussions with C. W. J. Beenakker, E. McCann, V. Fal'ko, B. Györfy, B. Nikolić, and A. Piróth. This work is supported by E. C. Contract No. MRTN-CT-2003-504574.

-
- ¹ K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, Y. Zhang, S. V. Dubonos, I. V. Grigorieva, and A. A. Firsov, *Science* **306**, 666 (2004); K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, *Nature* **438**, 197 (2005).
 - ² Y. Zhang, J. P. Small, M. E. S. Amori, and P. Kim, *Phys. Rev. Lett.* **94**, 176803 (2005); Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, *Nature* **438**, 201 (2005).
 - ³ M. I. Katsnelson, K. S. Novoselov, A. K. Geim, *cond-mat/0604323*.
 - ⁴ C. W. J. Beenakker, *cond-mat/0604594*.
 - ⁵ S. Bhattacharjee and K. Sengupta, *cond-mat/0607489*.
 - ⁶ M. Titov and C. W. J. Beenakker, *Phys. Rev. B* **74**, 041401 (2006).
 - ⁷ V. Lukose, R. Shankar, and G. Baskaran, *cond-mat/0603594*.
 - ⁸ B. Trauzettel, Ya. M. Blanter, A. F. Morpurgo, *cond-mat/0606505*.
 - ⁹ D. V. Khveshchenko, *cond-mat/0607174*.
 - ¹⁰ P. G. Silvestrov and K. B. Efetov, *cond-mat/0606620*.
 - ¹¹ V. V. Cheianov and V. I. Fal'ko, *Phys. Rev. B* **74**, 041403 (2006).
 - ¹² Cs. Tóke, P. E. Lammert, V. H. Crespi, and J. K. Jain, *cond-mat/0606461*.
 - ¹³ Y. Yao, F. Ye, X.-L. Qi, S.-C. Zhang, and Z. Fang, *cond-mat/0606350*.
 - ¹⁴ J. W. McClure, *Phys. Rev.* **104**, 666 (1956); D. P. DiVincenzo and E. J. Mele, *Phys. Rev. B* **29**, 1685 (1984).
 - ¹⁵ E. Fradkin, *Phys. Rev. B* **63**, 3263 (1986).
 - ¹⁶ P. A. Lee, *Phys. Rev. Lett.* **71**, 1887 (1993).
 - ¹⁷ E. V. Gorbar, V. P. Gusynin, V. A. Miransky, and I. A. Shovkovy, *Phys. Rev. B* **66**, 045108 (2002).
 - ¹⁸ A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, *Phys. Rev. B* **50**, 7526 (1994).
 - ¹⁹ V. P. Gusynin and S. G. Sharapov, *Phys. Rev. Lett.* **95**, 146801 (2005).
 - ²⁰ N. M. R. Peres, F. Guinea, and A. H. Castro Neto, *Phys. Rev. B* **73**, 125411 (2006).
 - ²¹ M. I. Katsnelson, *Eur. J. Phys B* **51**, 157 (2006).
 - ²² J. Tworzydło, B. Trauzettel, M. Titov, A. Rycerz, C.W.J. Beenakker, *Phys. Rev. Lett.* **96**, 246802 (2006).
 - ²³ K. Ziegler, *cond-mat/0604537*.
 - ²⁴ L. A. Falkovsky and A. A. Varlamov, *cond-mat/0606800*.
 - ²⁵ K. Nomura and A. H. MacDonald, *cond-mat/0606589*.
 - ²⁶ V. P. Gusynin and S. G. Sharapov, *cond-mat/0512157*.
 - ²⁷ K. S. Novoselov, E. McCann, S. V. Morozov, V. I. Fal'ko, M. I. Katsnelson, U. Zeitler, D. Jiang, F. Schedin, A. K. Geim, *Nature Physics* **2**, 177 (2006).
 - ²⁸ E. McCann and V. I. Fal'ko, *Phys. Rev. Lett.* **96**, 086805 (2006).
 - ²⁹ M. Koshino and T. Ando, *cond-mat/0606166*.
 - ³⁰ M. I. Katsnelson, *cond-mat/0606611*.
 - ³¹ A. J. McKane and M. Stone, *Annals of Physics* **131**, 36 (1981).
 - ³² B. A. Bernevig, *Phys. Rev. B* **71**, 073201 (2005).
 - ³³ J. Cserti and Gy. Dávid, *cond-mat/0604526*.
 - ³⁴ F. Guinea, A. H. Castro Neto, and N. M. R. Peres, *Phys. Rev. B* **73**, 245426 (2006); J. Nilsson, A. H. Castro Neto, N. M. R. Peres, and F. Guinea, *Phys. Rev. B* **73**, 214418 (2006); B. Partoens, and F. M. Peeters, *Phys. Rev. B* **74**, 075404 (2006).
 - ³⁵ V. P. Gusynin, S. G. Sharapov, and J. P. Carbotte, *cond-mat/0603267*.